HADRONIC ENERGY FLOW AND INCLUSIVE CHARGED PARTICLE PRODUCTION AT HERA

Stuart Mark Robertson

Thesis submitted for the degree of
Doctor of Philosophy

School of Physics and Space Research
Faculty of Science
University of Birmingham

September 1995

Synopsis

The energy flow in hard photoproduction interactions has been studied using data collected by the H1 detector at HERA during the 1993 data taking period. The production of jets and the energy flow about the resulting jet axis have been studied using a cone jet algorithm. The increased energy flow about the jet axis for jets reconstructed in the proton direction, compared with predictions of Monte Carlo event generators incorporating leading order pQCD calculations together with initial and final state radiation, is illustrated. The description of the energy flow has been found to be improved when multiple interactions models have been included.

The presence of multiple interactions has been further explored by studying charged particle production in the region $2.0 < \eta < 2.5$. To facilitate this the performance of the H1 Forward Tracker has been studied in detail, and the corrected track multiplicity measured in this region. This has further illustrated the inadequate description of photoproduction provided by leading order Monte Carlo generators and the improved description provided by the inclusion of multiple interaction models.

Cross sections are presented for the inclusive photoproduction of charged particles in the kinematical region $2.0 < \eta < 2.5$ and $p_T > 1.5$GeV/c. The transverse momentum distribution is found to be well described by a QCD inspired power law fit, and is found to be harder than observed in $p\bar{p}$ interactions at comparable centre of mass energies, $\sqrt{s_{\gamma p}} \approx \sqrt{s_{p\bar{p}}} \approx 200\,\mathrm{GeV}$.

This work was supported financially by the Particle Physics and Astronomy Research Council

Acknowledgements

I would like to thank Professor John Dowell for giving me the opportunity to work for the High Energy Physics Department, and for his help and guidance as my supervisor.

I am also grateful to the members of the H1 photoproduction working group for providing an interesting and relaxed environment for analysis during my stay at DESY. In particular, I am indebted to Gerhard Knies for his guidance and enthusiasm, and Mark Burton for his help with the Forward Tracker (good luck with the writing up).

I would also like to thank all the members of staff at Birmingham. In particular, I would like to thank Ian Kenyon and John Garvey for their help with getting to grips with HERA physics and H1. I would also like to thank Laurie Lowe for answering all my computer questions.

During my studentship, I have been fortunate to meet many interesting and lively people, who have made my time at Birmingham and DESY very enjoyable. I am grateful to Lee West, Paul Newman, Paul Sutton, Tim Nicholls, Kirstee Hewitt, Vicky Hudgson, Tanwir Ahmed and Hywel Phillips for their help as fellow H1'ers. Thanks Lee and Rachel for your culinary experiments which have prevented me from fading away during the dark days of writing up. I hope you both have an excellent time out at Hamburg. I am also grateful to you all for the many late night beer sessions in Hamburg.

I would also like to thank all the OPAL and OMEGA students at Birmingham. In particular I would like to thank Andy, Jim and Steve for their help as fellow newbies all those years ago, and for providing me with countless opportunities to test my liver. Thanks also to Chris and Rich, I look forward to joining you and Jim down in Harrow to ruin the reputation of particle physics in the real world forever. I must also mention Paul Davies, Keith Norman, Dave Rigby, Mark Venables, Dave Rees, Dave Evans, James Bloomer and Andy Bell for providing such an entertaining working environment.

Thanks also to everyone I met in Hamburg, in particular Dave Kant, John Lomas, Pete Bispham, Mark Burton, Paul Sutton and Dave Milstead. I must also mention Rob Akers, for forcing me to get drunk even when I didn't want to.

Finally I would like to thank my parents and family for their support (not least of all financially!) during my studies.

Contents

List of Figures

List of Tables

Chapter 1

Introduction

Elementary particle physics is concerned with the search for a description of the fundamental structure of matter and the forces of nature. During the last couple of decades, dramatic progress has been made in this field with the advent of huge particle accelerators enabling physicists to study matter at the most fundamental level. From the confusing profusion of experimental data on particles and phenomena, a theory known as the **Standard Model** has been developed which has been able to predict successfully many of the recent precise experimental results in terms of a small number of fundamental particles and interactions.

1.1 The Standard Model

In the Standard Model, matter is believed to be constructed from a small number of fundamental, point-like fermions. The forces between these particles, with the exception of gravity, have been successfully described by the exchange of fundamental gauge bosons within the framework of **Gauge Field Theories**.

Two classes of fundamental fermions exist, known as quarks and leptons. These are arranged in three quark and lepton families. Each lepton family consists of a negatively charged particle and a corresponding neutral, massless (or very light) neutrino. A total of six types or flavours of quarks are believed to exist. The existence of the five lightest quarks has been well established by experimental measurements.

	CHARGE	GENERATIONS I	II	III
QUARKS	+2/3	up	charm	top
	-1/3)	down	strange	bottom
LEPTONS	-1	e	μ	τ
	0	ν_e	ν_μ	ν_τ

Table 1.1: The three generations of fermions within the Standard Model.

Recently, experimental evidence of the existence of the sixth quark, known as the 'top' quark, has been observed by the CDF [1] and D0 [2] collaborations in proton-antiproton collisions at the Fermilab Tevatron accelerator.

The quarks are grouped into three families consisting of a quark with electric charge $+\frac{2}{3}$e (the *up (u), charm (c)*, and *top (t)* quarks) and a quark with electric charge $-\frac{1}{3}$e (the *down (d), strange (s)*, and *bottom (b)* quarks). A 'generation' then consists of a family of quarks and the corresponding family of leptons, as illustrated in table 1.1. It is believed that only three generations of quarks and leptons exist. Excluding antiparticles, the Standard Model includes 12 fundamental fermions.

It is believed that all the observed, strongly interacting particles of nature (referred to as hadrons) consist of either 3 quarks (or anti-quarks), referred to as baryons, or a quark and an antiquark pair, referred to as mesons. This model of baryons and mesons (known as the static quark model) was very successful in describing observed hadronic spectroscopy.

The fundamental forces of nature are described in the Standard Model using the language of quantum field theory. At the energy scales of interest to particle physics gravity is in fact very weak and can be neglected. The remaining three forces that have been observed are electromagnetism, the weak nuclear force and finally the strong nuclear force. All the observed fundamental fermions experience the electromagnetic force (with the exception of the electrically neutral neutrinos) and the weak nuclear force. However only the quarks experience the strong nuclear force.

Electricity and magnetism were unified into a single electromagnetic force by Maxwell in 1860. The weak nuclear force was postulated in order to explain phenomena such as β-decay. Finally the strong nuclear force was postulated as being the force that holds the atomic nucleus together.

1.1.1 Quantum Electrodynamics

Electromagnetism was the first force to be described by an accurate quantum field theory known as **Quantum Electrodynamics** or QED. In QED the electromagnetic force is mediated by the exchange of a massless gauge boson known as the photon. This is one of the most successful theories ever developed and is an example of a gauge field theory since it is based around the idea of gauge symmetry.

Quantum electrodynamics is formulated within the Lagrangian framework, in which the energy-momentum dynamics of a system is described by a Lagrangian density (often referred to as the Lagrangian). Quantum electrodynamics is based on the postulate that the Lagrangian should be invariant under an arbitrary change in the phase of the fermionic field at any point in space and time. This is known as a 'local gauge' transformation, as opposed to a 'global gauge' transformation in which a constant phase transformation is applied at all points in space and time. Whilst the initial Langrangian of a fermionic field is invariant under a global gauge transformation, the Lagrangian is not invariant under local transformations. In order to enforce local gauge invariance, additional terms must be introduced to the Lagrangian. One of the most important discoveries of particle physics is that the resulting terms introduced to the Langrangian correspond to a term equivalent to the Lagrangian describing the Maxwell equations of electromagnetism (the photon field), together with a term describing the interaction between the fermion and photon fields. Furthermore local gauge invariance also requires that the photon be massless.

The interaction term introduced to the Lagrangian by imposing local gauge invariance enables predictions to be made which may then be compared with experimental measurements. Calculations are performed using perturbation theory, in which each subsequent order of expansion in terms of the electromagnetic coupling constant, α, corresponds to one or more unique Feynman diagrams. Unfortunately, simple application of perturbation theory introduces divergent integrals into the calculation. However, an important property of QED is that it is 'renormalizable'. In this procedure the divergent integrals obtained are absorbed into a redefinition of the 'bare' lepton charges and masses, which are in any case arbitrary, so that the new definitions are equal to the observed physical values.

Another important consequence of QED is that the electromagnetic coupling constant is in fact not a constant and instead is dependent on the energy scale of the interaction. This can be understood in terms of the cloud of virtual electron-positron pairs that surround a bare electron. The cloud of electrons and positrons will tend to be polarised, screening the bare electron charge. The more penetrating the probe into the electron structure the larger the charge of the electron will appear to be. The coupling constant is therefore referred to as a 'running coupling constant'.

1.1.2 The Electro-Weak Interaction

After the success of the unification of electricity and magnetism, and the description of the subsequent electromagnetic force using QED, the Glashow-Weinberg-Salam model was proposed in order to describe the electromagnetic and weak interactions in a unified manner. In this model, the electromagnetic and weak interactions are considered to be aspects of the same force, and at very high energy scales have the same strength. However, as the weak interaction is mediated by the exchange of the massive $W^{\pm}$ and Z gauge bosons, the force appears weak at low energies because of its short range.

Experimentally, the charged current weak interaction was found to couple to left handed fermions only, whilst the neutral current interaction is known to couple to both left and right handed fermions. The parity violating nature of the charged current interactions is accounted for by treating the left and right-handed fermions differently. The left handed fermions are grouped into weak isospin doublets, and the right handed fermions are treated as weak isopin singlets. The electroweak interaction is then described by requiring invariance under the gauge transformation $SU(2)_L \times U(1)_Y$. The term $SU(2)_L$ refers to the gauge group of rotations in weak isospin space. The term $U(1)_Y$ corresponds to the gauge group of weak hypercharge (Y), where weak hypercharge, Y, is defined by

$$Q = T^3 + \frac{Y}{2}$$

where Q is the electric charge and T^3 is the third component of weak isospin, i.e. zero for isospin singlets and $\pm\frac{1}{2}$ for doublets. Imposing invariance of the Lagrangian under the above gauge transformation results in the introduction of four gauge fields.

A consequence of the above gauge invariance is that the observed bosons must be massless. However, the short range of the weak interactions indicates that in fact the $W^{\pm}$ and Z bosons are massive with a mass $\mathcal{O}(100\,\text{GeV})$. Furthermore, if a mass term is introduced to the Lagrangian, the resulting theory is no longer renormalizable. This problem has been solved by the 'Higgs mechanism', in which four Higgs fields are introduced to the Lagrangian which enables the possibility of having a 'spontaneously broken symmetry'. This means that the Lagrangian is still invariant under local $SU(2)_L \times U(1)_Y$ transformations, but in the ground state the invariance is broken enabling the gauge bosons to acquire mass. Combining three of the four Higgs fields with the weak bosons results in the massive $W^{\pm}$ and Z bosons, requiring the introduction of just one additional physical boson, known as the Higgs boson. A detailed discussion of electroweak physics and the Higgs mechanism is beyond the scope of this thesis.

1.1.3 Quantum Chromo-Dynamics (QCD)

Finally, the last observed fundamental force, the strong nuclear force, is described in the language of quantum field theory by **Quantum Chromo-Dynamics** or QCD. The strong interaction acts on colour charge, analogous to the electric charge of electromagnetism. The quarks are believed to exist in three 'colour' states, referred to as red, green and blue. Colour was first introduced to describe the wavefunction of particles such as the Δ^{++} particle for which the combined space, spin and flavour wavefunction is symmetric under the exchange of identical fermions. However the overall wavefunction is expected to be antisymmetric. This was resolved by introducing the colour degree of freedom, with the colour wavefunction being antisymmetric, restoring the required overall antisymmetric nature of the full wave function describing the Δ^{++} baryon.

QCD is based on the gauge transformation $SU(3)_C$ of colour. The requirement of gauge invariance under such transformations leads to the introduction of an octet of bi-coloured (i.e. carrying colour-anticolour quantum numbers) gauge bosons known as gluons. The fact that the resulting gluons themselves carry colour implies that the gluons also experience the strong interaction. This is referred to as the non-Abelian nature of QCD.

The non-Abelian nature of QCD has important consequences for calculations performed using perturbation theory. In analogy to QED, the strength of the strong interaction is described by the strong coupling constant, α_s, which again is a running coupling constant. However, due to the non-Abelian nature of QCD, the cloud of gluons surrounding a quark has the opposite effect to the cloud of electrons and positrons surrounding a bare electron. Instead of screening the colour charge of the quark, the effective colour charge of the quark increases with increasing distance (or equivalently decreasing four momentum transfer Q^2). This is believed to be the reason why free quarks have not been observed and instead quarks are always confined within colourless hadrons. However, as $Q^2 \to \infty$ the strong coupling constant decreases and the quarks appear to be weakly interacting. This is known as 'asymptotic freedom'. Consequently, QCD can only be solved using perturbation theory for high momentum transfers ($Q^2 > \mathcal{O}(\text{a few GeV}^2)$). As a result, though the basic interactions involved in many high energy scattering experiments can be calculated using perturbative QCD, the production of the final state hadrons seen in the detector cannot be calculated, and instead phenomenological models are required to describe the resulting hadronisation process.

Chapter 2

Physics at HERA

2.1 Introduction

Throughout the history of physics, scattering experiments have been crucial to the development of our understanding of the fundamental structure of matter and the forces of nature. Early scattering experiments by Rutherford using α particles led directly to the discovery of the nucleus. Later experiments by Hofstadter using electrons led to the measurement of the sizes of various nuclei and the observation of proton constituents within the helium nucleus[3]. Observations of point-like partons within the proton itself have resulted from the scattering of leptons (e, μ, ν) at various fixed target experiments [4]. The experiments H1 and ZEUS, using the HERA accelerator at DESY, are the latest in this chain of experiments and have the potential to extend our understanding of the structure of matter and the fundamental forces of nature still further.

2.2 Electron-Proton Scattering at HERA

At HERA energies, electron-proton (ep) scattering is dominated by the inelastic scattering process :

$$e^-(k) + p(P) \longrightarrow e^-(k') + X$$

where X is the hadronic final state. The four momentum transfer squared, Q^2, is defined as

$$Q^2 \equiv -q^2 = -(k - k^{'})^2 \tag{2.1}$$

where q is the four momentum of the virtual exchange particle. By convention, the above definition is used so that Q^2 is a positive quantity. Such interactions, in which the exchanged boson is either a virtual photon or Z^0, are referred to as 'Neutral Current' (NC) interactions.

Interactions in which a W^- boson is exchanged of the general form

$$e^- + p \longrightarrow \nu_e + X$$

are also possible and are referred to as 'Charged Current' (CC) interactions. Interactions involving the exchange of a Z^0 or W^- are only significant at high Q^2 as the cross section is suppressed by the factor

$$\frac{\sigma(Z^0, W^-)}{\sigma(\gamma)} \propto \left| \frac{Q^2}{Q^2 + M^2_{Z^0,W^-}} \right| \tag{2.2}$$

with respect to photon exchange.

High Q^2 ($Q^2 > \mathcal{O}(4\,\mathrm{GeV}^2)$) interactions are referred to as 'Deep Inelastic Scattering' (DIS). Interactions in which the momentum transfer is small ($Q^2 < \mathcal{O}(1\,\mathrm{GeV}^2)$), mediated by the exchange of a quasi-real photon, are referred to as 'photoproduction'.

2.3 Deep-Inelastic Scattering

The kinematics of deep inelastic scattering can be described at any given centre of mass energy by two independent variables. An example could be the energy and scattering angle of the scattered lepton. However, it is convenient to work in terms of the Lorentz invariant quantities Q^2 and x, the Bjorken scaling variable, defined as

$$x \equiv Q^2/(2P.q). \tag{2.3}$$



Figure 2.1: A schematic representation of deep inelastic scattering in lowest order QCD and QED.

Additional useful variables are

$$y \equiv (P.q)/(P.k) \tag{2.4}$$

and

$$W^2 \equiv (q+P)^2 \tag{2.5}$$

the invariant mass of the hadronic final state. A schematic representation of a deep inelastic ep interaction in lowest order QCD and QED is shown in figure 2.1.

The Bjorken scaling variable, x, can be interpreted as the fraction of the proton's momentum carried by the struck quark in the infinite momentum frame of the proton [5]. The variable y can be interpreted as the fraction of the incident lepton's energy transferred to the proton, measured in the rest frame of the incident proton. The variables Q^2, x and y are related by the equation

$$Q^2 = sxy \tag{2.6}$$

where s is the squared centre of mass energy given by

$$s \equiv (k+P)^2 \simeq 4E_eE_p \tag{2.7}$$

and E_e and E_p are the electron and proton beam energies respectively.

At the energies available at the HERA accelerator it is possible to make measurements at much higher values of Q^2 (up to $\mathcal{O}(10^4)\,\mathrm{GeV}^2$) and, equally importantly, at much lower values of x than otherwise possible at fixed target experiments.

2.3.1 The NC Deep Inelastic Cross Section

The total differential neutral current deep inelastic cross section can be expressed as :

$$\frac{d^2\sigma_{tot}}{dxdQ^2} = \frac{d^2\sigma_B}{dxdQ^2} + \frac{\alpha}{2\pi}\frac{d^2\sigma_1}{dxdQ^2} + \mathcal{O}(\alpha^2) \tag{2.8}$$

where the first term represents the Born cross section and the second and third terms represent the electro-weak radiative corrections to the Born cross section. In order to determine the Born cross section, the effect of radiative corrections must be accounted for. The largest contribution to the difference between the measured cross section and the Born cross section is due to initial state radiation, in which the incoming electron emits a real photon before interacting with the proton. For such events, the kinematics determined at the electron vertex are not the true kinematics of the ep interaction.

The Born cross section can be expressed in terms of two independent functions, $F_1^P(x,Q^2)$ and $F_2^P(x,Q^2)$, known as structure functions, by the equation

$$\frac{d^2\sigma_B}{dxdQ^2} = \frac{4\pi\alpha^2}{xQ^4}\left[y^2 x F_1^P(x,Q^2) + (1-y)F_2^P(x,Q^2)\right] \tag{2.9}$$

2.3.2 The Quark-Parton Model

In the quark-parton model, deep inelastic scattering is viewed as a two stage process. In the first stage, the virtual exchanged photon interacts with a quark within the proton, which over the short time scale of the interaction is assumed to be free. In the second stage of the process the outgoing partons hadronise to produce a 'current jet' of hadrons due to the struck quark, and a proton remnant.

The Born cross section can then be expressed in terms of the cross section for electron-quark scattering by the equation

$$\frac{d^2\sigma_B}{dxdQ^2} = \frac{4\pi\alpha^2}{xQ^4}\sum_q \int_0^1 d\xi f_q(\xi) e_q^2 \frac{x}{2}\left[1+(1-y)^2\right]\delta(x-\xi). \tag{2.10}$$

Equating equations 2.9 and 2.10 the following relationships are obtained :

$$F_2^P = 2xF_1^P \tag{2.11}$$

$$F_2^P = \sum_q e_q^2 x f_q(x) \tag{2.12}$$

It can be seen that, in the quark parton model, the structure function F_2^P is only dependent on the Bjorken scaling variable, x, and not on Q^2. This is known as Bjorken Scaling[6]. Equation 2.11 is known as the Callan-Gross relationship.

Within the framework of the quark parton model, if the total momentum of the proton was carried by the proton's quark constituents, then the momentum sum rule

$$\sum_i \int_0^1 dx x f_i(x) = 1 \tag{2.13}$$

would be expected to be true, where the sum is over all quark flavours. However, early measurements at SLAC [7] found that this relationship was not true with only approximately 50% of the total momentum of the proton being carried by the quark constituents. This was the first direct evidence for the existence of gluons within the proton.

2.3.3 Scaling Violations

In DIS measurements at previous fixed target experiments, Bjorken Scaling has been observed to be approximately true for $x \approx 0.15$. However, significant scaling violations have been observed at both lower and higher values of x. Such scaling violations can be predicted using the QCD improved quark-parton model. In QCD it is possible for a quark to radiate a gluon and for the gluon subsequently to split into a $q\bar{q}$ pair. At low Q^2 this structure is not resolved, however as the Q^2 of the probe is increased the structure is observed. Consequently, at high Q^2 there is an increased probability to observe a parton with a low value of x and conversely a reduced probability to observe a parton with a high value of x. Therefore, at fixed x the proton structure function F_2 will increase (decrease) with increasing Q^2 for low (high) x respectively. This leads to the 'Q^2 contraction' of the measured structure function.

The scaling violations can be predicted within the framework of perturbative QCD (pQCD) using the Altarelli-Parisi evolution equations [8] in which the evolution of the quark distribution functions, $q_i(x, Q^2)$, are given by

$$\frac{dq_i(x,Q^2)}{d\log Q^2} = \frac{\alpha_s}{2\pi}\int_x^1 \frac{dz}{z}\left(q_i(z,Q^2)P_{qq}\left(\frac{x}{z}\right) + g(z,Q^2)P_{qg}\left(\frac{x}{z}\right)\right) \tag{2.14}$$

where $g(x, Q^2)$ is the gluon distribution function which evolves with Q^2 according to the equation

$$\frac{dg(x,Q^2)}{d\log Q^2} = \frac{\alpha_s}{2\pi}\int_x^1 \frac{dz}{z}\left(\sum_i q_i(z,Q^2)P_{gq}\left(\frac{x}{z}\right) + g(z,Q^2)P_{gg}\left(\frac{x}{z}\right)\right). \tag{2.15}$$

The splitting function $P_{qq}(\frac{x}{z})$ gives the probability that a quark with momentum fraction z ($z > x$) will radiate a gluon giving a quark with momentum fraction x and can be calculated from pQCD. Equation 2.14 also includes a term due to the possibility of a gluon splitting into a $q\bar{q}$ pair (P_{qg}). Similarly, the evolution equation for the gluon distribution function includes two terms corresponding to the probability for a quark to radiate a gluon (P_{gq}) and the probability for a gluon to radiate a gluon (P_{gg}) due to the triple gluon vertex.

The Altarelli-Parisi equations therefore give a set of coupled differential equations, which given the quark distribution functions at a given $Q^2 = Q_0^2$ can be solved to determine the evolution of the quark distribution function for all $Q^2 > Q_0^2$. Generally, an analytical form of the parton distribution function at the input scale is used, the free parameters of which are then fitted by comparing with the measured structure functions.

2.4 Photoproduction

Though the primary interest at HERA may be with high Q^2 deep inelastic scattering, the ep cross section at HERA is in fact dominated by low Q^2 photoproduction events in which the interaction is mediated by a quasi-real photon. Photoproduction is however very interesting in its own right and offers a rich potential for physics analysis. Photoproduction offers an ideal testing ground for QCD in both the perturbative and non-pertubative regimes. In particular, the study of high p_T photoproduction with the production of jets enables the determination of the partonic structure of the photon.

For photoproduction events, the electron vertex can be treated to a good approximation by the 'equivalent photon approximation' (EPA). In this approximation, the cross section is factorised into the flux factor to describe the probability for a photon to be emitted at the electron vertex multiplied by the γp cross-section for real photons. Using the Weizacker-Williams approximation [9] the incident electron beam can be considered as a source of real photons with a flux given by

$$\frac{d^2 f}{dy dQ^2} = \frac{\alpha}{2\pi Q^2}\left(\frac{1+(1-y)^2}{y} - \frac{2m_e^2 y}{Q^2}\right). \tag{2.16}$$

A discussion of the limitations of the Weizacker-Williams approximation and photon flux calculations in general can be found in [10, 11] and [12]. Consequently the measured photoproduction cross section can be related to the total γp cross-section using the relationship

$$\frac{d^2 \sigma_{ep}}{dy dQ^2} = \frac{\alpha}{2\pi Q^2}\left(\frac{1+(1-y)^2}{y} - \frac{2m_e^2 y}{Q^2}\right)\sigma_{\gamma p}^{tot}(ys). \tag{2.17}$$

Several different mechanisms contribute to photoproduction at HERA energies. To first approximation, a real photon is a fundamental pointlike particle. However, through its direct coupling to quarks, a real photon has the possibility to split into a $q\bar{q}$ pair. The $\gamma \rightarrow q\bar{q}$ fluctuation may have varying virtualities, which can be characterised by the common p_T^2 of the q and $\bar{q}$ with respect to the photon direction. When the virtuality is small a vector meson wave function description may be a good approximation. This is the basis behind the vector dominance model (VDM) [13, 14] in which the photon couples to a hadronic vector particle with the same quantum numbers as the photon, which in turn interacts strongly with the proton.

In the VDM model it is postulated that the photon wavefunction is a superposition of a bare photon component, $\mid \gamma_B >$, and a hadronic component,

$\sqrt{\alpha} \mid h >$,

$$| \gamma >= \sqrt{Z_3} \mid \gamma_B > +c\sqrt{\alpha} \mid h > \tag{2.18}$$

where $Z_3 = 1 - c^2\alpha$ assures proper normalisation. At fixed target energies, the general features of photoproduction interactions have been found to be remarkably similar to hadronic interactions, for example πN interactions. This implies that the bare photon interactions can only account for a small fraction of the total photoproduction cross section: the smallness of the $\sqrt{\alpha} \mid h >$ component is more than compensated by the intrinsically much larger hN cross section. In the VDM model, interactions involving the bare photon component are neglected. Furthermore it is postulated that the three vector mesons ρ^0,ω and ϕ are the sole hadronic constituents of the photon, based on their copious photoproduction. The inclusion of heavier constituents is referred to as generalised vector dominance (GVD) [15].

The VDM model has been successfully applied in describing photoproduction at fixed target energies and has illustrated the similarity between photoproduction and hadron-hadron interactions. In hadronic physics, interactions can be separated into several different event classes such as elastic, diffractive and non-diffractive events. This classification can be extended naturally to the VDM model.

The elastic process $V + p \rightarrow V + p$ generalises to the 'VDM elastic' process[1]

$$\gamma + p \rightarrow V + p \tag{2.19}$$

The VDM single and double diffractive processes are correspondingly

$$\gamma + p \rightarrow V + X \tag{2.20}$$

$$\gamma + p \rightarrow X + p \tag{2.21}$$

$$\gamma + p \rightarrow X_1 + X_2 \tag{2.22}$$

where for case 2.22 the event is characterised by a large rapidity gap between the photon and proton dissociative systems. The remaining events,

$$\gamma + p \rightarrow X \tag{2.23}$$

[1]The truly elastic (Compton) process $\gamma + p \rightarrow \gamma + p$ is $\mathcal{O}(\alpha^2)$ and can safely be neglected.

where the hadronic system X does not contain any significant rapidity gaps, are referred to as non-diffractive events.

The non-diffractive photoproduction cross-section is dominated by 'soft' events leading to low p_T particle production. However, the observation of the production of high p_T particles and the production of jets in photoproduction has clearly shown that there is also a hard component to the non-diffractive cross section. Two components contribute to this hard component of the cross-section. Firstly the photon, as the gauge particle mediating the electromagnetic interaction, can interact directly with a parton from the proton. This class of events is known as 'direct photoproduction'. The second contribution occurs when the photon is resolved into a hadronic system, and a hard scattering occurs between a parton from the photon and a parton from the proton. Such interactions are called 'resolved photoproduction'.

The direct contribution to the non-diffractive cross-section can be calculated using perturbative QCD. Resolved photoproduction includes a component due to the VDM contribution. However, the pointlike coupling of the photon to quarks also leads to to a contribution where the photon splits into a $q\bar{q}$ pair, at relatively high virtuality, without forming a bound state. This additional mechanism is referred to as the 'anomalous' contribution to the photoproduction cross-section [16, 17, 18]. Figure 2.2 shows the typical direct, VDM and anomalous contributions to the hard non-diffractive cross section.

2.4.1 The Photon Structure Function

The above interpretation of the structure of the photon and its interactions leads to the introduction of parton distribution functions within the photon. The hadronic structure of the photon has been studied at e^+e^- colliders in $\gamma^*\gamma$ interactions. The real γ is radiated by an electron scattered at a small angle, whilst the highly virtual photon γ^* is 'radiated' by an electron scattered at a large angle in the detector. In analogy to the measurement of the structure of the proton in deep inelastic ep scattering, the resulting differential cross section can be expressed in terms of the photon structure functions F_1^γ and F_2^γ [19]. The resulting differential cross section



Figure 2.2: A schematic representation of the typical (a) direct, (b) VDM and (c) anomalous contributions to the hard non-diffractive photoproduction cross section.

for the interaction $e\gamma \rightarrow eX$ where X is any hadronic final state is then given by

$$\frac{d^2\sigma(e\gamma \rightarrow eX)}{dxdQ^2} = \frac{4\pi\alpha^2}{xQ^4}\left[y^2xF_1^\gamma(x,Q^2) + (1-y)F_2^\gamma(x,Q^2)\right]. \qquad (2.24)$$

The evolution of the photon structure functions from a given input scale, Q_0^2, to higher Q^2 can then be calculated in a similar manner to F_2^P using the Altarelli-Parisi equation with the inclusion of a term to account for the anomalous contribution to the photon structure function [20].

The situation is consequently very similar to the case of nucleonic parton densities. However, the determination of the input distributions is much more difficult in the case of the photon. In particular, no momentum sum rules apply for the photon and it is consequently difficult to derive reliable information on the gluon content of the photon. Furthermore, the $e\gamma \rightarrow eX$ event rate in e^+e^- interactions is quite small; most recent measurements are based typically on around 1,000 events at $Q^2 \simeq 5\,\text{Gev}^2$ [21, 22, 23]. This can be compared with millions of events in, for example, deep-inelastic μN scattering.

Several parameterisations of the parton distribution functions of the photon exist. In particular, three leading order (LO) parton distribution sets have been produced by Levy, Abramowicz and Charchula (LACI-III) [24], and leading order and next to leading order (NLO) parton distribution sets have been produced by Gluck, Reya and Vogt (GRV-LO and GRV-NLO) [25].

The LAC parton distributions are based on the QCD evolution of input distributions at $Q_0^2 = 4\,\text{GeV}^2$ (LACI and LACII) and $Q_0^2 = 1\,\text{GeV}^2$ (LACIII) respectively. An analytical form of the input distribution at Q_0^2 was assumed with a total of 12 free parameters. The free parameters were then determined by a fit to all experimental data on F_2^γ above the initial value, Q_0^2. The set LACII was produced with one less free parameter. The resulting parton distribution functions illustrate the poor constraint on the gluon content of the proton provided by data on F_2^γ. In particular the LACIII parameterisation predicts a very hard gluon content, with a maximum at $x \simeq 0.9$, whilst the LACI and LACII parameterisations predict very soft gluon distributions, rising very steeply at low x.

The GRV parton distribution functions (GRV-LO and GRV-NLO) are based on the QCD evolution of input distributions at a very low Q_0^2 ($0.25\,\mathrm{GeV}^2$ in LO, $0.3\,\mathrm{GeV}^2$ in NLO). The approach is based on the same philosophy as used in the previous fits of protonic and pionic parton densities by the same authors. In the case of the proton, it was initially hoped that the complex structure of the proton could be generated dynamically from a valence quark distribution at the chosen input scale. However, whilst the gluon density was found to evolve sufficiently to carry approximately half of the proton's momentum at Q^2 of a few GeV^2, the gluon distribution was found to be too soft in shape. Consequently 'valence-like' gluon and sea quark densities had to be introduced even at the input scale. Similarly the pionic input distributions also included a 'valence-like' gluon density, though no sea quarks were introduced at the input scale.

The GRV parton distribution functions for the photon were subsequently obtained by using input distributions of the form

$$\vec{q}^{\,\gamma}_{0,GRV}(x) = \kappa \frac{4\pi\alpha_{em}}{f_\rho^2} \vec{q}^{\,\pi}_{0,GRV}(x) \tag{2.25}$$

where $\vec{q}^{\,\pi}_{0,GRV}(x)$ are the corresponding input distribution functions used for case of the pion. The use of the above input distributions is motivated by VDM considerations, where f_ρ^2 determines the $\gamma \to \rho$ transition probability. The constant κ is introduced to allow for the contribution of heavier vector mesons, and is a free parameter in the subsequent fit to the available data. Equation 2.25 is an assumption which must be tested experimentally; in particular it is not clear that the parton densities in a pion are similar to those of a vector meson, or that pQCD is applicable at such low scales.

A general review of the theoretical uncertainties in the current knowledge of the photon structure functions can be found in [18].

2.4.2 The Total Photoproduction Cross Section

The total photoproduction cross-section has been measured by both the H1 and ZEUS collaborations at centre of mass energies of $\sqrt{s} \simeq 200$ GeV. Using the data collected in the 1992 data taking period, the total photoproduction cross section was measured by the H1 collaboration as

$$\sigma_{tot}^{\gamma p} = 159 \pm 6.6(stat) \pm 20(sys)\mu b \tag{2.26}$$

where the first error is statistical and the second error is systematic [26]. The measurement is consistent with the total cross section determined by the ZEUS collaboration [27]. However, in order to determine the cross section a model dependence is introduced as the acceptance corrections are sensitive to the expected contribution to the cross section of the various event classes discussed above.

Using a data sample collected during a short dedicated running period in 1994, the H1 collaboration has attempted to reduce the systematic error in the above measurement by measuring the elastic and diffractive contributions to the total cross section[28]. The total cross section was then determined as

$$\sigma_{tot}^{\gamma p} = 165 \pm 2(stat) \pm 11(sys)\mu b. \tag{2.27}$$

The resulting cross section is only then dependent on the assumed rate of the double diffractive contribution to the total cross section. The measurement shows the typical increase in the total cross section at high energies compared with previous measurements using fixed target experiments at $\sqrt{s} \leq 18$ GeV, as observed in hadronic interaction cross sections.

Predictions of the dependence of the total cross section in photoproduction are based on two main approaches: (i) Regge-type model predictions and (ii) QCD-based predictions in 'mini-jet' models.

In Regge theory [29] the high-energy behaviour of a two-body amplitude due to the exchange of a Regge pole is given by

$$A(s,t) \sim s^{\alpha(t)} \tag{2.28}$$

where $\alpha(t)$ is the relevant Regge trajectory. Using the optical theorem the total cross section is related to the imaginary part of the forward scattering amplitude, giving

$$\sigma_{tot} \sim s^{\alpha(0)-1} \tag{2.29}$$

where $\alpha(0)$ is the intercept of the Regge trajectory

The Regge pole carrying the quantum numbers of the vacuum is called the Pomeron, while all other poles are referred to as Reggeons. The total cross section for the hadronic interaction $A + B \to X$ can then be described by the general form

$$\sigma_{tot}^{AB} = X^{AB} s^{\alpha_P(0)-1} + Y^{AB} s^{\alpha_R(0)-1}. \tag{2.30}$$

The constants $\alpha_P(0)$ and $\alpha_R(0)$ are believed to be universal, with a recent fit to measured hadronic cross sections [30] finding $\alpha_P(0) = 1.0808$ and $\alpha_R(0) = 0.5475$. The coefficients X^{AB} and Y^{AB} are process dependent, and must be determined by a fit to previous cross section measurements for the process under consideration. A recent fit to photoproduction data below 20 GeV gives a predicted cross section at 200 GeV of 160 μb [17], which is consistent with the measured cross section determined by the H1 and ZEUS collaborations.

In the second class of predictions on the total photoproduction cross section, the increase in the total cross section in hadronic interactions at high centre of mass energies is interpreted as due to the increase in the cross section for the production of two (mini-)jets in the framework of pQCD [31, 32]. In LO pQCD, parton parton scattering leads to the production of two outgoing jets. The scale that is used to determine the point at which perturbative calculations are applicable is usually taken as the transverse momentum squared (p_T^2) of the outgoing partons. The di-jet cross section is given by pQCD as

$$\frac{d\sigma_{\gamma p}^{jet}}{dp_T^2} = \sum_{i,j,k} \int dx_\gamma \int dx_p \int d\hat{t} f_i^\gamma(x_\gamma, \mu^2) f_j^p(x_p, \mu^2) \frac{d\hat{\sigma}_{ij}^k}{d\hat{t}} \delta\left(p_T^2 - \frac{\hat{t}\hat{u}}{\hat{s}}\right) \tag{2.31}$$

where $f_i^{\gamma,p}$ are the parton distribution functions and $\hat{\sigma}$ is the partonic scattering cross section. Consequently the total di-jet cross section above a given $p_T = p_{T,min}$ is then given by

$$\sigma_{\gamma p}^{jet}(p_{T,min}) = \int_{p_{T,min}^2}^{s/4} \frac{d\sigma^{jet}}{dp_T^2} dp_T^2. \tag{2.32}$$

It is reasonable that an effective $p_{T,min}$ cut should be applied since the incoming hadrons are colour neutral objects and when the p_T of the exchanged gluon (say) is small and the transverse wavelength correspondingly large, the gluon can no longer resolve the individual colour charges, and the effective coupling is decreased. This mechanism cannot be predicted by pQCD calculations, which are always performed assuming the scattering of free partons, but is also not in contradiction with pQCD.

If $\sqrt{s} \gg p_{T,min}$ equation 2.32 receives its dominant contribution from $x_{\gamma,p} \ll 1$. The parton density functions can then be approximated by a simple power law, $f_i^\gamma, f_j^p \propto x^{-J}$ where the power J is expected to lie roughly in the range $1.2 \leq J \leq 1.5$. The total di-jet therefore cross section scales at high (asymptotic) energies like [33]

$$\sigma_{\gamma p}^{jet}(p_{T,min}) \propto \frac{1}{p_{T,min}^2}\left(\frac{s}{4p_{T,min}^2}\right)^{J-1} \log \frac{s}{4p_{T,min}^2}. \tag{2.33}$$

The cross section will therefore grow much faster than the total γp cross section, which only grows $\propto \log^2 s$ (Froissart bound [34]). Eventually the jet cross section will therefore exceed the total γp cross section.

This apparent problem is solved by the observation that pQCD is performed assuming free partons, whilst the incoming hadrons can be viewed as beams of partons. Consequently a multiplicity factor can be introduced. Since each partonic scattering leads to a pair of (mini-)jets the following relationship is obtained :

$$\sigma_{ab}^{jet} = \langle n_{jetpairs}\rangle\, \sigma_{ab}^{inel}. \tag{2.34}$$

This is known as multiple scattering and can be included within the eikonal formalism. In this approach the eikonal $\chi(b,s)$ is introduced, where χ is identified as the mean density of hard scatters for collisions at impact parameter b. χ is assumed to be of the form

$$\chi(b,s) = \frac{A(b)\sigma^{jet}(s)}{P_{had}} \tag{2.35}$$

where $A(b)$ describes the overlap of partons within the colliding particles in impact parameter space and P_{had} is the probability that both initial particles are in a hadronic state when they interact. The overlap function $A(b)$ is usually taken as a convolution of the electromagnetic form factors of the incident particles and P_{had} is expected to be $\sim 4\pi\alpha_{em}/f_\rho^2(\simeq 1/300)$. If the multiple partonic interactions are assumed to be independent at a fixed impact parameter, then the probability that no hard scattering should occur is given by Poisson statistics as $e^{-\chi(b,s)}$. The total



Figure 2.3: An example of multiple parton scattering in γp interactions.

cross section is then given by

$$\sigma_{tot}^{pQCD} = P_{had} \int_0^\infty \pi db^2 \left[1 - \exp(-\chi(b,s))\right] \tag{2.36}$$

which has manifestly taken into account the possibility of more than one hard scatter per interaction, illustrated in figure 2.3

In realistic analyses of the total photoproduction cross section a non-perturbative contribution to the eikonal is introduced:

$$\chi(b,s) = \frac{A(b)\left[\sigma^{soft} + \sigma^{jet}(s)\right]}{P_{had}} \tag{2.37}$$

with the soft cross section often parameterised as

$$\sigma^{soft}(s) = \sigma_0 + \frac{\sigma_1}{\sqrt{s}} \tag{2.38}$$

where σ_0 and σ_1 are constants. Therefore including $p_{T,min}$ three free parameters (beyond those describing high p_T jet production) are available to fit the total photoproduction cross section.

Extreme predictions from mini-jet models are excluded by the recent measurements of the total photoproduction cross secion. However, minijet jet predictions could be consistent with the measured cross section provided a $p_{T,min}$ value of about 2 GeV/c is chosen.

Jets with a transverse momentum of $\mathcal{O}$(1-3 GeV/c) are referred to as mini-jets. Recently the TOPAZ and AMY collaborations unambiguously observed [35] mini-jets in $\gamma\gamma$ collisions at TRISTAN. These data, as well as other data on multi-hadron production in $\gamma\gamma$ scattering [36] imply that partonic collisions with transverse momenta in the GeV range occur at the rate predicted by pQCD. A detailed review of the uncertainties in recent predictions of the total photoproduction cross section based on mini-jet models can be can be found in [37, 38].

It should be emphasised that it is not possible to determine the presence of multiple parton-parton interactions by an analysis of the total cross section alone. In particular, the measured cross section can equally well be described by Regge parameterisations as discussed above.

2.4.3 Jet Production

The production of high p_T jets offers the largest cross-section of all hard γp scattering reactions. Consequently, the production of jets in photoproduction has been the focus of much of the early analysis by the H1 [39, 40, 41, 42] and ZEUS [43, 44, 45, 46] collaborations at HERA. In LO pQCD, hard photoproduction leads to the production of two outgoing partons leading to the production of two jets at the experimental level.

The kinematics of a typical resolved photoproduction event is shown in figure 2.4. The variables x_γ and x_P are the momentum fraction of the parton from the photon and proton respectively. For events with two well reconstructed jets it is possible to reconstruct x_γ and x_P from the measured transverse energy, E_T, and pseudorapidity, η, of the reconstructed jets using the equations

$$x_\gamma = \frac{E_T^1 exp(-\eta_1) + E_T^2 exp(-\eta_1)}{2E_\gamma} \tag{2.39}$$

$$x_P = \frac{E_T^1 exp(\eta_1) + E_T^2 exp(\eta_1)}{2E_P}. \tag{2.40}$$



Figure 2.4: The kinematics of a typical LO resolved photoproduction event. The production of two outgoing partons leads to the observation of two hadronic jets in the event, characterised by their transverse energy, E_T, and pseudorapidity, η.

As described in the previous section it is possible for multiple interactions to occur in resolved photoproduction. The presence of multiple interactions should lead to observable features in hard photoproduction events. In particular the presence of additional mini-jets or soft activity in the event leads to an increased 'jet pedestal', where the term jet pedestal refers to the energy flow away from the jet axis. The increased jet pedestal should also contribute additional energy to the jet, leading to an increased inclusive jet rate. The situation is however complicated as other effects such as initial and final state radiation and fragmentation also contribute to the energy flow in such events. A detailed discussion of jet production and the evidence for multiple interactions is presented in chapter 6.

The description of the energy flow in hard photoproduction is also important in order to enable the observed distribution of x_γ to be unfolded back to the true distribution in order to study the photon structure function. The gluon density in the photon has been studied by the H1 collaboration [41] by subtracting the contribution due to direct photoproduction (concentrated at high x_γ) and resolved



Figure 2.5: The gluon density of the photon divided by the fine structure constant α=1/137 (data: circles) at the scale $< p_T >^2 = 75\,\mathrm{GeV}^2$. For comparison the GRV-LO (full) and the LACI (dashed) and the LACIII (dotted) parameterisations are shown. From [41]

photoproduction events initiated by a quark from the photon from the unfolded x_γ distribution. The gluon density of the photon has then been measured at a scale of $< p_T >^2=75\,\mathrm{GeV}^2$ and is shown in figure 2.5 compared with predictions from the GRV-LO, LACI and LACIII parameterisations. It can be seen that the LACIII prediction of a very hard gluon density in the photon is ruled out by the measurement. The LACI distribution, giving a very steeply rising gluon density at low x_γ, is also not favoured, whilst the GRV-LO predictions give reasonable agreement with the data. For this reason, in the analysis presented in this thesis the GRV-LO parton distribution functions for the photon have been used in order to describe photoproduction interactions using Monte Carlo generators.

The above determination of the gluon density within the photon has been obtained by applying a jet pedestal subtraction method, as a full description of the energy flow in such events has not yet been achieved. A quite large model dependence is therefore introduced into the result. In order to improve the determination of the parton distribution functions in the photon from HERA data, a thorough description of the energy flow in such events is required. Consequently, the study of the presence of multiple interactions in photoproduction at HERA energies is essential for practical purposes, in addition to the theoretical interest in multiple interactions. This has been the motivation for much of the analysis presented in this thesis.

Chapter 3

The H1 Detector

3.1 The HERA accelerator

The HERA accelerator at DESY in Hamburg is the first electron-proton storage ring to be constructed. It is designed to accelerate 210 bunches of electrons (or positrons) and 210 bunches of protons to 30 GeV and 820 GeV respectively in two independent storage rings housed in a tunnel of circumference 6.3 km. The electron and proton bunches are arranged to intersect at four interaction points, two of which house the general purpose detectors H1 and ZEUS.

The layout of the HERA accelerator and pre-accelerators in shown in figure 3.1. The design of the electron and proton storage rings differ due to the respective limiting factors. For electrons the beam energy is limited by synchrotron radiation. However, for protons the effect of synchrotron radiation is negligible, and consequently the proton beam energy is limited by the magnetic field attainable in the bending magnets. Therefore, the electron ring uses conventional magnets operating at room temperature whilst the proton ring uses superconducting magnets. The proton beam is accelerated by a series of warm 52 and 208 MHz RF cavities in order to achieve the design beam energy and the electron beam is accelerated by a series of 82 warm cavities and 16 superconducting cavities.

Luminosity was first achieved at HERA during 1992. For the first year of running HERA operated with 10 electron and proton bunches, with the bunch crossing



Figure 3.1: The layout of the HERA accelerator and pre-accelerators.

interval set at the design value of 96 ns. However only 9 of the electron and proton bunches were collided. The 10^{th} bunch in both the electron and proton beams were separated, such they did not have a colliding partner. The non-colliding bunches, referred to as pilot bunches, enable beam induced backgrounds to be studied. The 1992 data taking period yielded an integrated luminosity of approximately $30\,\mathrm{nb}^{-1}$. For the 1993 data taking period several improvements enabled HERA to operate with 84 colliding electron and proton bunches, and 6 proton and 10 electron pilot bunches. A total integrated luminosity of $880\,\mathrm{nb}^{-1}$ was delivered.

3.2 The H1 Detector

In order to realise the potential of the HERA accelerator, the H1 detector has to satisfy certain basic criteria. The identification and measurement of electrons plays a crucial role in the physics at HERA, and consequently the H1 detector must be able to facilitate this. The detector must have good electromagnetic and hadronic energy resolution in order to study inclusive energy flow measurements both in DIS and photoproduction. It is important that the detector is as hermetic as possible to facilitate the measurement of missing transverse energy, in order to select and study charged current DIS events. Muon identification and energy resolution must be excellent in order to allow the study of interactions that produce heavy flavours, as well as to allow the search for new particles.

An accurate determination of the luminosity is required, and this must also include the facility to study photoproduction via the tagging of low angle electrons produced in low Q^2 interactions. Finally a beam crossing occurs at the H1 detector every 96 ns, and consequently sophisticated triggers are required in order to select physics events from the large numbers of beam-gas and beam-wall background interactions, and to distinguish between different physics classes.

The layout of the H1 detector is shown in figure 3.2. A right-handed Cartesian coordinate system (x, y, z) is used, where z is the direction of the proton beam, y is vertical and the origin is at the nominal interaction point. A spherical polar coordinate system (r, θ, ϕ) is defined in the conventional way with reference to the Cartesian coordinate system, such the $\theta = 0°$ is the proton direction. The regions $\theta < 90°$ and $\theta > 90°$ are referred to as the 'forward' and 'backward' region respectively.

The main components of the H1 detector, as operational for the 1993 data taking period, are :

- A tracking detector consisting of separate central $(15° < \theta < 165°)$ and forward $(7° < \theta < 25°)$ components together with a backward MWPC in the electron direction.

- A calorimeter system consisting of the main electromagnetic and hadronic LAr calorimeters, supplemented by a backward electromagnetic calorimeter (BEMC) in the electron direction, a plug calorimeter to detect small angle hadronic energy in the very forward direction, and limited streamer tubes (LST) in the iron return yoke to measure hadronic energy that has leaked out of the main calorimeter system.



Figure 3.2: The H1 detector (3D view).

- A superconducting coil and cryostat outside the hadronic calorimeter in order to provide a magnetic field of approximately 1.15 Tesla over the central region.

- A set of iron plates surrounding the superconducting coil to act as a return yoke for the magnetic field. The return yoke is instrumented both to act as a tail-catcher for hadronic energy and also to provide muon identification and tracking.

- A forward muon spectrometer to supplement the muon identification in the forward region.

- A luminosity monitoring system consisting of small angle electron and photon detectors in the electron direction. The small angle electron tagger is also used as a trigger for low-Q^2 physics.

The above components will be discussed in more detail in the following sections, with particular emphasis on those components of relevance to the analysis presented in this thesis. A more detailed description of the H1 detector can be found in [47].

3.3 The H1 Tracking Detectors

Due to the typical event topology at HERA, the tracking system is divided into two main components, the Central Tracking Detector (CTD), covering the angular range $15° < \theta < 165°$, and the Forward Tracking Detector (FTD), covering the angular range $7° < \theta < 25°$. The H1 tracking system is then completed by a backward multiwire proportional chamber (BPC) in front of the backward electromagnetic calorimeter. The BPC covers the angular range $155.5° < \theta < 174.5°$. A side view of the H1 tracking system is shown in figure 3.3. A superconducting coil provides a uniform magnetic field inside the tracking region of approximately 1.15 Tesla parallel to the beam direction.

Both the central and forward tracking detectors consist of a series of drift chambers and multiwire proportional chambers (MWPC). The drift chambers are designed to give accurate measurement of the trajectory of charged particles, enabling



Figure 3.3: Side view of the H1 tracking detectors.

a determination of the particle momentum. This is achieved by a system of anode (sense) and cathode wires, which are used to set up a uniform drift field within the chamber, with the exception of the region very close (<1 mm) to the sense wires where the field increases very rapidly. A charged particle transversing the chamber causes ionization along its path. The resulting electrons drift towards the sense wires and in the region of increasing electric field an avalanche of electrons is produced. This causes a detectable signal on the sense wires. Readout and timing at the ends of the wires enables the trajectory of the charged particle to be accurately reconstructed.

The multiwire proportional chambers are used in event triggering. The proportional chambers are made up of finely spaced wires, such that a charged particle experiences a very high initial electric field, independent of the particle's trajectory. The resulting pulses are detected very quickly and enable a fast determination of the approximate trajectory of the charged particle. This enables a fast trigger decision to be made. The spatial resolution of the MWPCs are limited by the spacing of the sense wires.

3.3.1 The Central Tracking Detectors

The functions of accurate $r\phi$- and z-measurement are separated in the central tracking detector. Accurate $r\phi$-measurement is provided by the Central Jet Chamber (CJC) [48] of which there are two rings. The CJC also provides moderate z-measurement by charge division; however precise z-measurement is instead provided by z-drift chambers both inside and outside the CJC. Finally two sets of multiwire proportional chambers provide a central track trigger [49]. Starting from the interaction region a charged particle is detected in the Central Tracking Detector by the inner MWPCs (CIP), the inner z-chamber (CIZ), the inner jet chamber (CJC1), the outer z-chamber (COZ), the outer MWPCs (COP) and finally the outer jet chamber (CJC2).

The Central Jet Chambers are JADE type jet chambers, with cells tilted by approximately 30° with respect to the radial direction and wires running parallel to the beam axis (see figure 3.4). The chamber has an inner radius of 200 mm, an outer radius of 795 mm and a length of 2640 mm centred about the nominal interaction point. A spatial resolution of $\sigma_{r\phi} \simeq 350\,\mu$m and $\sigma_z \simeq$ 5cm has been achieved [47], which compares well with the design goals.

The inner and outer z-chambers are drift chambers with wires stretched in a polygon around a cylinder. They consist of 16 azimuthal sectors and 60 cells in the z-direction. Using this arrangement a z-resolution of $\sigma \simeq 300\,\mu$m has been achieved [47]. The inner and outer z chambers are combined in a package with the inner and outer MWPCs respectively. The inner z-chambers and inner MWPCs are designed to fit into a cylinder with an inner radius of 15 cm, an outer radius of 20 cm and a total length of 249 cm. The outer z-chambers and the outer MWPCs lie between the two CJC chambers, with an inner radius of 45.5 cm and an outer radius of 49 cm.



Figure 3.4: Cell structure of the Central Jet Chamber

3.3.2 The Forward Tracking Detectors

The H1 Forward Tracking Detector (FTD) [50] consists of three identical sub-units, known as supermodules, numbered from 0 to 2 in increasing z. Each supermodule, when seen from the direction of the incoming proton, consists of a planar chamber module, followed by a MWPC, then a layer of transition radiator material, and finally a radial drift chamber. The FTD overs the angular range between 7° and 25° with a sensitive region of $134 < z < 254$ cm and $15 < R < 79$ cm. The orientation of the chambers is designed to maximise the number of sense wires passed by a particle and so increase the number of hits per track. The layout of the H1 forward tracker is illustrated in figure 3.5

Each planar chamber module consists of three planar drift chambers orientated



Figure 3.5: Side view of the H1 Forward Tracker.

at $0°, +60°$ and $-60°$ to the vertical respectively. Each chamber consists of four wire planes in the z direction. The sense wires in each drift chamber are only read out at one end of the wires, and hence no information is available concerning the coordinate of each hit along the wires. However the separate orientations enable the trajectory of a particle passing through the tracker to be reconstructed. The structure of the planar drift chambers is illustrated in figure 3.6. The methods used to reconstruct a track in the planar chambers will be discussed in more detail in chapter 5.

The radial drift chambers consist of planes of sense wires radiating out from the beam-pipe, such that the maximum drift length for each wire increases with increasing distance from the beam pipe. This enables an accurate measurement of the azimuthal angle, together with a moderate measurement of the radial coordinate using charge division. The radial chambers are also used to detect the X-rays from the transition radiators. Each radial chamber has 48 cells in ϕ equipped with 12 sense wires stacked in z. The layout of the radial drift chambers is illustrated in figure 3.7



Figure 3.6: The Planar Drift Chambers.

A more detailed description of the forward tracker and the reconstruction of tracks in this region is given in chapter 5.

3.3.3 The Backward MWPC

The H1 tracking system is completed by a multi-wire proportional chamber situated in front of the backward electromagnetic calorimeter. This consists of four planes of parallel wires, each rotated by 45° with respect to each other. The backward MWPC completes the trigger system for the tracking detector and also provides additional space points for small angle tracks pointing towards the backward electromagnetic calorimeter in order to improve the identification and measurement of electrons for low-Q^2 DIS events.



Figure 3.7: The Radial Drift Chambers.

3.4 The H1 Calorimeters

The H1 tracking system is complemented by the H1 calorimeter system which is designed to measure the total energy of both charged and neutral particles produced in the interaction. The calorimeter system consists of the main Liquid Argon Calorimeter (LAr) covering the angular range $4^\circ < \theta < 153^\circ$, the Backward Electromagnetic Calorimeter (BEMC) covering the angular range $151^\circ < \theta < 177^\circ$, the Plug Calorimeter in the very forward direction ($0.3^\circ < \theta < 3^\circ$) and the Tail Catcher (TC) covering the angular range $6^\circ < \theta < 172^\circ$.

The H1 calorimeters are all based on the same principle of layers of passive absorption material interleaved with layers of instrumented active regions. A particle passing through the absorption material produces a shower of particles. The shower particles produced lose energy by ionisation or excitation of the active medium. By measuring the ionisation produced in the active regions the development of the shower can be periodically sampled. Consequently, the total energy of the incident particle can be measured and the longitudinal and transverse profile of the shower can be determined.

In electromagnetic calorimeters, energetic photons and electrons produce a shower of lower energy photons and electrons due to Bremsstrahlung and pair production processes. The development of an electromagnetic shower in a given absorption material is characterised by the radiation length, X_0, the mean longitudinal length over which an electron loses all but 1/e of its energy.

A strongly interacting particle passing through an absorption material loses energy through inelastic nuclear collisions producing lower energy hadrons which undergo further inelastic collisions. Hadronic showers take longer to develop than electromagnetic showers, and for any given absorption material the longitudinal development of the hadronic shower is given by the nuclear interaction or absorption length, λ.

The LAr calorimeter [51] consists of an electromagnetic section with lead absorbers, corresponding to a depth of between 20 and 30 radiation lengths, and a hadronic section with steel absorbers. The total depth of the LAr calorimeter varies between 4.5 and 8 hadronic interaction lengths. The containment of the LAr calorimeter is shown in figure 3.8 in terms of radiation lengths $\mathrm{X_0}$ and absorption lengths λ for the electromagnetic and hadronic components respectively. The calorimeter is highly segmented in both sections, with a total of around 45000 cells. The root mean square electronic noise per channel typically corresponds to 10 to 30 MeV.

Test beam studies of LAr calorimeter modules have demonstrated energy resolutions of $\sigma(E)/E \simeq 0.12/\sqrt{E} \oplus 0.01$ with E in GeV for electrons [52] and $\sigma(E)/E \simeq$